


AP 330-1 Anaphylaxis Emergency Plan

 (Student Name)

This Person has a potentially life-threatening allergy (anaphylaxis) to:



Check the appropriate box(es):

<input type="checkbox"/>	Peanut	<input type="checkbox"/>	Tree Nuts
<input type="checkbox"/>	Egg	<input type="checkbox"/>	Milk
<input type="checkbox"/>	Insect Sting	<input type="checkbox"/>	Medication: _____
<input type="checkbox"/>	Latex	<input type="checkbox"/>	Other: _____

Food: The key to preventing an anaphylactic emergency is absolute avoidance of the allergen. People with food allergies should not share food or eat unmarked/bulk foods or products with a 'may contain' warning.

Epinephrine Auto-Injector: Expiry Date _____

Dosage: EpiPen® Jr 0.15 mg EpiPen® 0.30 mg

Location of Auto-Injector(s): _____

(or student will carry his/her own medication)

Asthmatic: Person is at greater risk. If person is having a reaction and has difficulty breathing, give epinephrine auto-injector before asthma medication.

A person having an anaphylactic reaction might have ANY of these signs and symptoms:

- **Skin:** hives, swelling, itching, warmth, redness, rash.
- **Respiratory (breathing):** wheezing, shortness of breath, throat tightness, cough, hoarse voice, chest pain/tightness, nasal congestion or hay fever-like symptoms (runny itchy nose and watery eyes, sneezing), trouble swallowing.
- **Gastrointestinal (stomach):** nausea, pain/cramps, vomiting, diarrhea.
- **Cardiovascular (heart):** pale/blue colour, weak pulse, passing out, dizzy/lightheaded, shock.
- **Other:** anxiety, feeling of 'impending doom', headache.

Early recognition of symptoms and immediate treatment could save a person's life.

Act quickly. The first signs of a reaction can be mild, but symptoms can get worse very quickly.

- 1. Give epinephrine auto-injector** (e.g. EpiPen® or Twinject™) at the first sign of a reaction occurring in conjunction with a known or suspected contact with allergen. Give a second dose in 5 to 10 minutes or sooner **IF** the reaction continues or worsens.
- 2. Call 911.** Tell them someone is having a life-threatening allergic reaction. Ask them to send an ambulance immediately.
- 3. Go to the nearest hospital,** even if symptoms are mild or have stopped. Stay in the hospital for an appropriate period of observation, generally 4 hours, but at the discretion of the ER physician. The reaction could return.
- 4. Call contact person.** (see info on following page)

Emergency Contact Information:

Name	Relationship	Home Phone	Work Phone	Cell Phone

The undersigned patient, parent or guardian authorizes any adult to administer epinephrine to the above-named person in the event of an anaphylactic reaction as described above. This protocol has been recommended by the patient's physician, and has been reviewed with the patient/child and the parent/guardian.

Patient/Parent/Guardian Signature

Date

Physician Signature

Date